

PAVEL KOLÁŘ, PaedDr, PhD¹ • JAN ŠULC, MD, PhD² • MARTIN KYNČL, MD³ • JAN ŠANDA, Ing³ • ONDŘEJ ČAKRT, MSc²
ROSS ANDEL, PhD⁴ • KATHRYN KUMAGAI, DPT⁵ • ALENA KOBESOVÁ, MD, PhD²

Postural Function of the Diaphragm in Persons With and Without Chronic Low Back Pain

Despite the high prevalence of low back pain in the population, options regarding effective treatment strategies are still limited, possibly due to the lack of knowledge of the underlying mechanisms.⁹ Trunk stabilization and postural trunk control may play an important role in the etiology of low back pain.⁹ In turn, the function of the diaphragm may affect how the trunk is stabilized, especially during postural activity.^{11,15,22} Various studies have shown that the pelvic girdle and lumbar spine are reflexively stabilized and

braced prior to the initiation of extremity movements.^{3,13,17,22} The central nervous system must be able to anticipate movement and stabilize the entire core musculature automatically to provide a stable base from which the muscles performing the movement can pull.

Trunk bracing maintains all spinal segments in a biomechanically neutral position during the course of any movement. Segmental movement (eg, hip joint movement) is therefore related to the synergistic activity of the spinal extensors and all the muscles modulating intra-abdominal pressure (ie, abdominal muscles, the diaphragm, and the pelvic floor). The diaphragm is the muscle that contributes the intra-abdominal pressure modulation and plays an important role in spinal stability.^{7,16,18,27,31}

Insufficient function and poor coordination of postural, or stabilizing, muscles are considered to be important etiological factors in spinal disorders associated with low back pain, such as deformational spondyloarthrosis (with or without spinal disc herniation), spinal disc protrusion, and/or spondylolisthesis.^{5,12,21,25} However, a study designed specifically to test the dynamics of the diaphragm in chronic spinal disorders is lacking. We aimed

● **STUDY DESIGN:** A case-control study.

● **OBJECTIVES:** To examine the function of the diaphragm during postural limb activities in patients with chronic low back pain and healthy controls.

● **BACKGROUND:** Abnormal stabilizing function of the diaphragm may be an etiological factor in spinal disorders. However, a study designed specifically to test the dynamics of the diaphragm in chronic spinal disorders is lacking.

● **METHODS:** Eighteen patients with chronic low back pain due to chronic overloading, as ascertained via clinical assessment and magnetic resonance imaging, and 29 healthy subjects were examined. Both groups presented with normal pulmonary function test results. A dynamic magnetic resonance imaging system and specialized spirometric readings were used with subjects in the supine position. Measurements during tidal breathing (TB) and isometric flexion of the upper and lower extremities against external resistance with TB were performed. Standard pulmonary

function tests, including respiratory muscle drive (PI_{max} and PE_{max}), were also assessed.

● **RESULTS:** Using multivariate analysis of covariance, smaller diaphragm excursions and higher diaphragm position were found in the patient group ($P < .05$) during the upper extremity TB and lower extremity TB conditions. Maximum changes were found in costal and middle points of the diaphragm. A 1-way analysis of covariance showed a steeper slope in the middle-posterior diaphragm in the patient group both in the upper extremity TB and lower extremity TB conditions ($P < .05$).

● **CONCLUSION:** Patients with chronic low back pain appear to have both abnormal position and a steeper slope of the diaphragm, which may contribute to the etiology of the disorder. *J Orthop Sports Phys Ther* 2012;42(4):352-362, Epub 21 December 2011. doi:10.2519/jospt.2012.3830

● **KEY WORDS:** dynamic magnetic resonance imaging, lung function, spinal disorders, stabilizing function, thorax

¹Associate Professor, Department of Rehabilitation and Sport Medicine, Second Medical Faculty, Charles University and University Hospital Motol, Prague, Czech Republic.

²Lecturer, Department of Rehabilitation and Sport Medicine, Second Medical Faculty, Charles University and University Hospital Motol, Prague, Czech Republic. ³Lecturer, Department of Radiology, Second Medical Faculty, Charles University and University Hospital Motol, Prague, Czech Republic. ⁴Associate Professor, School of Aging Studies, University of South Florida, Tampa, FL; International Clinical Research Center, St Anne's University Hospital Brno, Brno, Czech Republic. ⁵Associate Professor, Department of Physical Therapy, Azusa Pacific University, Azusa, CA. This study was supported by the foundation Movement without Help, Prague, Czech Republic. This study was approved by the Second Medical Faculty, Ethical Committee, Charles University and University Hospital Motol, Prague, Czech Republic. Address correspondence to Dr Alena Kobesová, Department of Rehabilitation and Sport Medicine, Second Medical Faculty, Charles University and University Hospital Motol, V Úvalu 84, Prague 5, 15006, Czech Republic. E-mail: alenamudr@me.com

to examine diaphragm excursions and inspiratory/expiratory positions during normal tidal breathing (TB) and during postural tasks in patients with chronic low back pain and healthy volunteers. We hypothesized that the diaphragm excursions would be reduced in the patient group and that the inspiratory and expiratory positions of the diaphragm during postural tasks would be more cranial, reflecting abnormal diaphragm function. In addition, we expected that the recruitment manner of diaphragm sections during inspiration would show an altered contraction pattern in the patient group compared to healthy controls, indicating abnormal coordination. We assessed diaphragm positions using dynamic magnetic resonance imaging (MRI), with synchronized respiratory assessment during normal TB and postural limb activities, and analyzed respiratory muscle drive (maximum inspiratory/expiratory occlusion pressures).

METHODS

Participants

PARTICIPANT RECRUITMENT occurred in 2 phases. All participants were recruited at the University Hospital Motol in Prague, Czech Republic, by a single clinician (P.K.). In the first phase, a sample of 30 healthy participants was recruited as part of a previously published study of the stabilizing function of the diaphragm.²⁷ One participant was excluded due to insufficient data required for the present study. In the second phase, available volunteers with chronic low back pain due to vertebral disorders were assessed ($n = 18$).

The inclusion criteria of the patient group included (a) low back pain that was not due to a specific injury and that could be classified as chronic (lasting at least 6 months), (b) a lack of systemic disease that would contribute to low back pain or morphological changes, and (c) observation of morphological changes of chronic character in the lumbar spine due to chronic overloading (morphol-

ogy of the spinal disorder determined by MRI). Specifically, 7 patients were diagnosed with spondylosis and spinal stenosis, 2 patients with spondylosis and spondylolisthesis, 5 patients with spondylosis and disk hernia, and 4 patients with failed back surgery syndrome (patients operated for advanced spondylosis, spinal stenosis, and disk hernia not due to an injury). None of these conditions were a result of a spine or pelvic traumatic injury, or a chronic systemic or respiratory disease. Rather, we determined via clinical and MRI assessment that the morphological findings were consistent with abnormal stress to the structural elements of the lumbar spine. None of the patients suffered from chronic respiratory disease or such symptoms at the time of the study. Finally, participants who underwent a surgery for low back pain, listing injury as the primary cause, and patients in whom injury or a cause other than overuse or misuse contributed substantially to morphological changes in the lumbar spine were excluded.

The healthy participants (controls) included 4 males (14%) and 25 females (86%), with a mean \pm SD age of 29 ± 7 years and body mass index (BMI) of 22.3 ± 2.5 kg/m² (height, 168 ± 8 cm; weight, 64 ± 8 kg). The 18 patients with chronic low back pain included 11 males (61%) and 7 females (39%), with a mean age \pm SD of 48.6 ± 13 years and BMI of 26.8 ± 3.6 kg/m² (height, 177 ± 11 cm; weight, 84 ± 16 kg).

Procedures

This study was approved by The Institutional Ethical Committee. All participants underwent an in-person interview to ensure that they met the inclusion criteria of the study. All testing procedures were thoroughly explained to the participants with a detailed description of the dynamic MRI and spirometry assessments. All subjects reported that they understood the test procedures and gave their informed consent. Subjects were also instructed to fast at least 4 hours before each assessment procedure.

Participants were evaluated by dynamic MRI, with simultaneous respiratory recordings. Diaphragm activity, measured by excursion of the diaphragm, was evaluated by dynamic MRI, with subjects in the supine position, their heads supported 5 cm above the MRI plinth. Volumetric changes during the breathing cycle were recorded with a specially designed spirometer and specialized computer software. The subjects wore nose clips to prevent any air exchange through the nostrils. A mouthpiece connected to a pneumotachograph was placed in the subject's mouth, and the subjects were allowed to practice normal breathing through the mouthpiece. After the subjects were trained in normal breathing with the mouthpiece for 2 minutes, measurements were taken during TB at rest and again with isometric limb contractions of the upper and lower extremities. To ensure consistency during the testing procedures, the same physical therapist (P.K.) performed all assessments. Data collection time was 20 seconds in each condition per subject to record standard MRI measurements together with the respiratory readings.

Diaphragm activity was assessed under the following conditions (**FIGURE 1**). In all conditions, the subject was in the supine position with arms and legs relaxed along the torso. The subject was instructed to breathe normally throughout the assessment. The measurements of diaphragm movement and respiratory readings were recorded throughout the 20-second data collection period.

Tidal Breathing After the initial synchronization between respiratory and MRI recordings, simultaneous synchronized respiratory and MRI recordings were taken.

Isometric Flexion of Upper Extremity (UE) The clinician placed his hands on the dorsal surface of the subject's forearms, which were at rest. The subject was then instructed to keep the elbows straight and flex both shoulders, attempting to lift the arms against the clinician's resistance, maintaining an isometric con-




FIGURE 1. Schematic illustration of postural task performance during MRI assessment. (A) Initial resting supine condition. (B) Isometric flexion of lower extremity against the clinician's resistance: the arrow marked "F" corresponds with the direction of the clinician's resistance. The blue arrow represents the direction of required movement of the subject. (C) Isometric flexion of upper extremity against the clinician's resistance corresponding with the MRI assessment utilized in this study. Abbreviation: MRI, magnetic resonance imaging.

cranial diaphragm positions during TB in a healthy control (**FIGURE 2A**) and a patient with chronic low back pain (**FIGURE 2B**).

The diaphragm excursion images were converted to binary images to calculate their area in pixels. The bottom edge of the diaphragm excursion represents the most caudal baseline diaphragm position during inspiration. The top edge of the diaphragm excursion represents the diaphragm in its most cranial position during expiration. Successive images with the next highest pixel count were analyzed in order as the excursion of the diaphragm changed during the breathing cycle.

The next analysis was completed on the subtracted maximal crescent area of each image, where the horizontal, anterior/posterior alignment (perpendicular to the body axis), used as baseline, was represented in the front (point A in **FIGURE 2C**) by the inferior ventral costal margin, midclavicular line, and in the back (point E in **FIGURE 2C**) by the thoracolumbar junction in the dorsal axillar line. Markers designating the baseline were placed on each subject's body (the total anterior/posterior distance was linked with the dotted line from point A to point E). The total horizontal distance was divided into 6 equal sections, demarcating 5 equidistant points, with point C marking the midpoint of the line from points A to E (**FIGURE 2C**). The upper and lower edges of diaphragm excursion were determined as average change from the baseline at each of the 3 middle points (B, C, and D) across all breaths taken within the 20-second interval. The distance at each point from the horizontal baseline was calculated to determine the difference in inspiratory position compared to the expiratory position of the diaphragm in millimeters (B1-B2, C1-C2, and D1-D2) (**FIGURE 2C**).

Synchronized Respiratory Measurements

Synchronized respiratory recordings and their processing were performed using the approach described previously.²⁷ Tidal volumes were recorded with

traction (**FIGURE 1C**). The muscle power generated by the subject corresponded to a grade 4 manual muscle test.²⁶

Isometric Flexion of Lower Extremity (LE) The clinician placed his hands on the anterior surface of the subject's thighs while the subject remained at rest. The subject was then instructed to perform bilateral hip flexion against the clinician's resistance, maintaining an isometric contraction (**FIGURE 1B**). The muscle power generated by the subject corresponded to a grade 4 manual muscle test.²⁶

All subjects were able to achieve grade 4 force, without substantial pain. As the clarity of MRI images is negated by involuntary and/or excessive movement, the inevitable movement caused by substantial pain during the performed postural activities would have resulted in images that could not be interpreted.

MRI Assessments

MRI scans and MRI analysis of diaphragm movement were performed using the approach described previously.²⁷ The diaphragm was imaged in the sagittal plane with the subject in supine, us-

ing a body coil. The imaging plane was placed sagittally in the axial topogram directed paravertebrally to the right, midway between the center of the vertebral body and the edge of the thoracic wall. Slice thickness was 33 mm. Sequence duration was 20 seconds. Each subject had 3 markers (10-mL syringes of water) affixed to the skin surface and placed at (1) the midclavicular line, level of the jugular opening, (2) the inferior ventral costal margin, midclavicular line, and (3) the thoracolumbar junction in the dorsal axillar line.

The MRI image files were converted to Analyze format with MRICro software. In each 20-second sequence, the baseline position of the diaphragm was determined for TB and postural activity conditions. The most caudal baseline position of the diaphragm was subtracted from the position of the other images in the sequence to determine the position changes of the diaphragm throughout the 20-second collection period. **FIGURES 2A** and **2B** provide examples of the crescent-shaped image of diaphragm excursion, contrasting the most caudal and




FIGURE 2. (A) Subtracted image of the diaphragm excursions in the most caudal (inspiratory) and cranial (expiratory) diaphragm positions during tidal breathing in a healthy control. (B) A subtracted image of the diaphragm excursions in the most caudal (inspiratory) and cranial (expiratory) diaphragm positions during tidal breathing in a patient with chronic low back pain. (C) Schematic description of 3 diaphragmatic points (B, C, and D) used for diaphragm excursion calculations. The following 6 distances (in mm) were obtained by measuring the distance between the horizontal baseline in both expiratory and inspiratory diaphragm positions. Diaphragm excursion points: B1 to D1 were derived from the inspiratory diaphragm positions obtained from MRI images; B2 to D2 were derived from expiratory diaphragm positions obtained from respective MRI images. The inspiratory diaphragm position is designated by points B1, C1, and D1. The expiratory diaphragm position is designated by points B2, C2, and D2. Total diaphragm excursion is designated by the distance from the lower to the upper curve along points B1 to B2, C1 to C2, and D1 to D2. Adapted from Kolar et al.²⁷

a spirometer (MasterScope Jaeger Version 4.67; Jaeger, VIASYS, Würzburg, Germany) by a specially designed pneumotachograph with a plastic iso-resistive membrane. This device allowed safe and reliable respiratory recording while in a strong magnetic field. A specialized reading and recording BreathRecorder software (J. Volejník, Kurka-Jaeger Servis, Ltd, Czech Republic) was developed for the purposes of this study. The flow signal measured the airflow, which was then converted and digitally integrated to yield the measurement of volume using an analog-to-digital converter and saved on a hard disk. Prior to respiratory measurements, every subject was familiarized with the mouthpiece in a supine position for a 2-minute period, during which no recordings were performed. The recording system was calibrated to each subject using a 1-L calibration pump prior to data collection.

The respiratory data were processed using Software Grapher (J. Volejník, Kurka-Jaeger Servis, Ltd, Czech Republic). From the 20 seconds of recorded data in each condition, 4 to 7 respiratory cycles were used to calculate the tidal volume.

Synchronization of Respiratory Measurements and MRI Sequence The respi-

ratory measurements were synchronized at the beginning of the 20-second MRI sequence within the initial 200 to 300 milliseconds by an electronic marker imprinted simultaneously on both recordings. The individually marked respiratory recordings were converted to DICOM format and synchronized with the dynamic MRI sequence of diaphragm movement images. The synchronized progression of the trace volume-time respiratory curve and the corresponding diaphragm movement were monitored using RADinfo Scan View System software (Radiology Information Systems, Inc, Sterling, VA).

Pulmonary Function Tests

Standardized spirometric recordings²⁹ of pulmonary function tests (PFTs) were performed on the same day for all subjects with a MasterScope Jaeger spirometer (Version 4.5; Jaeger, VIASYS, Würzburg, Germany), with a special module for the assessment of respiratory muscles (drive). This is a widely used method that has been described elsewhere.² In brief, the patients were instructed to maintain maximum inspiratory and expiratory pressure for at least 1.5 seconds, so that the maximum pressure sustained for 1 second could be

recorded. The measured airway opening pressure for both maximum static inspiratory (PI_{max}) and maximum static expiratory pressure (PE_{max}) indicates global respiratory muscle output.² All subjects were properly instructed and coached by an experienced technician during all PFTs. Proper procedures for quality assurance, based on the criteria of the American Thoracic Society,²⁹ were used for these measurements. The following spirometric parameters were measured: forced expiratory volume in 1 second (FEV_1), forced vital capacity (FVC), and FEV_1/FVC . Concomitantly, assessments of respiratory muscles were performed. The following parameters used for these measurements were PI_{max} and PE_{max} . Procedures and quality criteria of the American Thoracic Society were used for these measurements.² The PFT results were compared to established reference values³³ using a regression equation that included age, sex, and height, which correlated most strongly with the respective normative values, along with the published regression coefficients as predictors of the observed values. The results are presented as percentages of the predicted values. The standard deviation of the residual of the predicted values (ie, the difference between the observed and

TABLE

INSPIRATORY AND EXPIRATORY POSITIONS OF THE DIAPHRAGM (MM)

	Controls		Cases		Mean Difference	95% CI [†]
	Mean	SD*	Mean	SD*		
Tidal breathing, inspiration						
Point B [‡]	94.1	29.4	92.8	26.1	1.3	-15.3, 18.0
Point C [‡]	94.8	30.9	92.9	29.3	1.9	-16.3, 20.0
Point D [‡]	77.9	33.5	71.0	31.4	7.0	-12.6, 26.5
Tidal breathing, expiration						
Point B	119.1	23.4	115.6	23.3	3.5	-10.7, 17.7
Point C	127.8	24.0	125.8	23.6	2.0	-12.5, 16.4
Point D	118.2	24.3	112.8	23.8	5.4	-9.2, 20.0
Upper extremity resistance, inspiration						
Point B	90.1	29.7	123.3	50.4	-33.1	-60.1, -6.1
Point C	88.3	31.3	121.3	52.8	-33.0	-61.3, -4.7
Point D	67.6	34.9	89.6	50.9	-22.0	-49.9, 6.0
Upper extremity resistance, expiration						
Point B	119.8	23.2	126.5	24.5	-6.7	-21.3, 7.9
Point C	127.6	23.8	139.2	27.2	-11.7	-27.5, 4.2
Point D	116.2	25.1	130.8	34.8	-14.6	-33.9, 4.7
Lower extremity resistance, inspiration						
Point B	85.7	32.6	120.1	46.1	-34.3	-57.5, -11.2
Point C	81.7	34.4	115.7	47.2	-34.0	-60.2, -7.7
Point D	59.7	38.2	82.8	46.8	-23.1	-49.9, 2.1
Lower extremity resistance, expiration						
Point B	116.5	25.4	125.2	25.3	-8.6	-24.0, 6.7
Point C	122.1	25.7	136.5	29.1	-14.5	-30.8, 1.9
Point D	109.0	28.1	125.3	36.7	-16.3	-36.9, 4.4

Abbreviations: CI, confidence interval.
**Standard deviation from the mean.*
[†]*Differences in means: lower and upper limits for the 95% CI around the difference in means; a result is significant when the CI does not cross zero.*
[‡]*Points B, C, and D refer to the anterior, middle, and posterior parts of the diaphragm, as illustrated in FIGURE 2.*

predicted values) was used to assess normality of the PFT results.

Pulmonary function tests yielded a predicted mean ± SD FEV₁ of 105.4% ± 9.6%, FVC of 109.7% ± 12.0%, and FEV₁/FVC of 99.5% ± 8.3% for the healthy controls, and FEV₁ of 106.1% ± 14.2%, FVC of 113.8% ± 16.0%, and FEV₁/FVC of 95.3% ± 11.6% for the patients with chronic low back pain. All observed values fell within 1 standard deviation of the residual of the predicted values, deeming them within the normal range.³³

Statistical Analysis

All analyses were performed using commercial software SPSS Version 15 (SPSS Inc, Chicago, IL). First, we used multi-

variate analyses of covariance (MANCOVAs) within the general linear model statistical framework³⁵ to assess group differences in diaphragm function. The MANCOVA yields an overall (multivariate) main effect, commonly represented by the F value associated with Wilks' lambda criterion. This statistic reflects the covariate-adjusted effect across the dependent variables, while taking into account the common variance shared by the dependent variables. The MANCOVA also yields covariate-adjusted main effects linked to the individual dependent variables, which we interpreted when the multivariate effect was statistically significant.

Group (patient group versus control

group) served as the independent variable across the models. The dependent variables were diaphragm excursions at each point of measurement, estimated separately for the TB, upper extremity TB (UETB), and lower extremity TB (LETB) conditions. In subsequent analyses, we examined group differences in the individual inspiratory or expiratory diaphragm positions by entering as the dependent variables the 3 points of measurement on the diaphragm during the TB, UETB, or LETB conditions.

We also tested whether coordination of the individual sections of the diaphragm (the recruitment manner) would be reduced during inspiration in patients with chronic low back pain. We




FIGURE 3. (A) Inspiratory and expiratory positions of the diaphragm during tidal breathing for the patient and control groups. (B) Inspiratory and expiratory positions of the diaphragm during tidal breathing with isometric flexion of the upper extremity in the patient and control groups. (C) Inspiratory and expiratory positions of the diaphragm during tidal breathing with isometric flexion of the lower extremity in the patient and control groups.

calculated a slope of the middle-posterior diaphragm for each participant by subtracting values for the middle and posterior inspiratory positions and compared mean values between groups using a 1-way analysis of covariance (ANCOVA), in which the slope of the middle-posterior diaphragm was the dependent variable, group (patient versus control) was the independent variable, and age and gender were the covariates.

To test coordination between the diaphragm and other respiratory muscles, we calculated Pearson correlation coefficients for the relationship between

MRI-derived total diaphragm excursion, specified as the sum of diaphragm excursion at points B, C, and D for each postural task (TB, UETB, and LETB), and the 2 indices of respiratory muscle drive (PI_{max} and PE_{max}) separately in the control and patient groups.

Statistical significance was assessed at a 2-tailed .05 level across all analyses.

RESULTS

THE TABLE PRESENTS THE UNADJUSTED means and standard deviations for diaphragm position during the 3 tasks. **FIGURE 3** illustrates inspiratory and expiratory diaphragm positions during normal TB (**FIGURE 3A**), UETB (**FIGURE 3B**), and LETB (**FIGURE 3C**).

The groups differed in age, gender, and body mass index (BMI). In an independent *t* test, with equal variances not assumed, the patient group was on average older ($t[24] = 6.04, P < .001$) and had higher BMI ($t[27] = 4.70, P < .001$) than the control group. There were more women than men in the patient group ($\chi^2 = 11.40, P < .001$). Subsequently, we also found that age and BMI were highly correlated ($r = .75, P < .001$) and, in a 1-way analysis of covariance, BMI was not statistically different in the patient group compared to the control group when age was used as a covariate ($F_{45} = 1.07, P = .31$), indicating that the higher BMI in the patient group was attributable to the higher age in this group. To use a parsimonious model more amenable to the relatively small sample size, only age and gender were used as covariates in the MANCOVAs.

Diaphragm Excursions and Inspiratory/Expiratory Diaphragm Positions During TB

During TB without postural movement, the MANCOVAs yielded no significant multivariate effects for the differences between the patient and control groups in diaphragm excursions, as well as diaphragm inspiratory and expiratory positions ($P > .05$) (**FIGURE 3A**).

Diaphragm Excursions and Inspiratory/Expiratory Diaphragm Positions During Postural Activities

Differences between the patient and control groups emerged when postural tasks were applied during TB (**FIGURES 3B** and **3C**). For diaphragm excursions, the MANCOVA-based multivariate effects for group were significant in both the UETB ($F_{1,41} = 3.69, P = .020$) and the LETB ($F_{1,41} = 4.76, P = .006$) conditions, indicating that diaphragm excursions were significantly smaller in the patient group compared to the control group under both conditions. In the UETB condition, the significant overall effect for group was attributable primarily to the group differences at point B ($P = .016$), whereas the differences at points C ($P = .059$) and D ($P = .601$) were not significant. In the LETB condition, the smaller diaphragm excursion in the patient group existed at points B ($P < .001$), C ($P < .001$), and D ($P = .023$).

We then assessed group differences in inspiratory and expiratory diaphragm positions individually. During the UETB condition, the multivariate main effect for group across the 3 points of measurement (B, C, and D) was significant during inspiration ($F_{1,41} = 4.11, P = .012$), indicating a higher (more cranial) position of the diaphragm in the patient group. This effect could not be attributed to any single point on the diaphragm ($P > .30$), although the group differences in diaphragm position appeared more pronounced at points B and C (**FIGURE 3B**). The multivariate main effect for group was not significant during expiration ($P = .336$).

During the LETB condition, the multivariate main effect for group was again significant during inspiration ($F_{1,41} = 3.49, P = .024$), with the diaphragm positioned higher in the patient group than the control group. As in the UETB condition, no single point represented this difference in diaphragm inspiratory position, with the group difference slightly larger at points B ($P = .132$) and C ($P = .141$) than D ($P = .361$). The multivariate main effect for group during expiration

PE_{max} or between diaphragm excursion and PI_{max} in the TB or UETB condition. However, in the LETB condition, positive correlation between diaphragm excursion and PI_{max} was found in the patient group ($r = .50, P = .035$) but not in the control group ($r = -.021, P = .916$). These 2 correlations are illustrated in **FIGURE 5**. Fisher r -to- z transformation indicated that the difference between the 2 coefficients approached statistical significance ($P = .078$).

DISCUSSION

WE STUDIED DIFFERENCES IN DIA- phragmatic activity during TB with and without postural activities in a group of patients with chronic low back pain compared to healthy volunteers. We found that diaphragm excursions or inspiratory/expiratory positions did not vary across patients and controls during TB without postural tasks. However, reduced diaphragm movement emerged when isometric flexion against resistance of the upper or lower extremity was applied, pointing to the importance of postural tasks in the expression of abnormal diaphragm function. The results provide additional evidence for altered breathing patterns during strenuous and nonstrenuous activity in individuals with low back pain.^{24,30} The findings may support the notion that the strategies utilized by the central nervous system to control core stability are altered in the presence of painful syndromes.¹⁴

Perhaps the most clinically important finding of this study concerns the abnormal coordination of the diaphragm in the patient group during inspiration with postural tasks. This impairment was demonstrated by reduced movement of the diaphragm in the anterior and middle portion, while the posterior (crural) part moved in the same manner as in the control group. This pattern of diaphragmatic recruitment resulted in a steeper angle in the middle-posterior part of the diaphragm (**FIGURE 4**), which may exacerbate the symptomology of chronic




FIGURE 4. The slope of the middle-posterior diaphragm during inspiration in the upper extremity tidal breathing condition for patients with chronic low back pain (A) and healthy controls (B).

was not significant ($P = .40$), although a post hoc analysis indicated that this effect was significant ($P = .022$) when age and gender were not controlled.

The Recruitment Manner of the Diaphragm

In this step, we examined the hypothesis that diaphragm contraction during inspiration occurs more unevenly in the middle-posterior portion in the patient group. Given previous evidence and the angle variation illustrated in **FIGURE 3**, we tested whether poor coordination in the patient group was represented specifically by altered diaphragm contraction in the middle-posterior (crural) portion of the diaphragm, denoted as points C and D. The unadjusted means on which this analysis was based are presented in the **TABLE**. In an analysis of covariance exam-

ining group differences in the slope of the middle-posterior diaphragm controlling for age and gender, the main effect for group was not significant during the TB condition ($P > .05$). However, there was a significant main effect for group during both the UETB ($F_{1,43} = 10.07, P = .003$) and the LETB ($F_{1,43} = 5.49, P = .024$) conditions, indicating that the contraction of the diaphragm followed a substantially steeper recruitment pattern in the patient group relative to the control group. For better illustration of this situation, **FIGURE 4** shows the slope of the middle-posterior diaphragm during inspiration in the UETB condition.

Total Diaphragm Excursion and Respiratory Muscle Drive

No significant correlations emerged between total diaphragm excursion and

low back pain by increasing the anterior shear forces on the ventral region of the spinal column.

Although the role of the ligamentous and muscular system to stabilize the low back and decrease the shear forces during loading has been studied extensively,^{28,32} not much attention has been paid to the coordination across individual sections of the diaphragm. Additionally, while radiological evaluation or visualization of the diaphragm is routinely performed in clinical practice, to our knowledge, no research to date has been conducted in which diaphragm dynamics would be visualized in relation to postural tasks and low back pain. In the control group, normal diaphragmatic contraction performed during inspiration was characterized by a symmetric recruitment of all diaphragmatic parts, that is, points B, C, and D. This motion is directed downward toward the abdominal cavity. The physiologically domed (convex) contour of the diaphragm at the end of expiration becomes less domed, and thus the contour is more symmetrical at the end of inspiration.

By contrast, although the patient group demonstrated normal recruitment of the crural portion of the diaphragm, movement in the anterior and medial portions was limited in both postural tasks. Poor coordination of particular diaphragmatic parts in the patients (points B and C) resulted in an asymmetric diaphragmatic activation during inspiration, as demonstrated by a steeper slope of the crural part of the diaphragm. Evidently, limited motion of the costal part may result in a more domed inspiratory diaphragmatic position. Contraction of the diaphragm has been found to modulate intra-abdominal pressure and contribute to trunk stability.^{11,19,20} Although we did not measure intra-abdominal pressure directly, we suspect that in the patient group, abnormal position and recruitment of the diaphragm resulted in subsequent reduced intra-abdominal pressure conducive to low back pain.¹⁸ These findings are consistent with the hypothesis




FIGURE 5. Correlation between diaphragm excursion during lower extremity flexion and maximum static inspiratory pressure (PI_{max}) in healthy controls (A) and patients with chronic low back pain (B).

that abnormal postural activation of the diaphragm may serve as 1 underlying mechanism of chronic low back pain.⁹

Finally, during LE flexion, total diaphragm excursions and maximum inspiratory pressure were significantly correlated in the patient group, whereas the parallel correlation in the control group was practically zero. It appears

that respiratory muscles other than the diaphragm were more consistently recruited during TB when lower extremities were engaged in order to compensate for the relatively limited movement of the diaphragm, whereas the diaphragm was more central to breathing in healthy subjects. However, we cannot assume that respiratory volume would neces-

sarily be reduced in the patient group. Previous research suggests that inhaled lung volume may even be greater in patients with low back pain,¹⁰ although this research did not include any measurement of diaphragm movement. Finally, we found pulmonary function within a normal range for both groups, further suggesting that compensation for the limited diaphragm movement might have occurred. Taken together, we speculate that an altered mode of activation, indicative of poor activation of the diaphragm during TB with postural tasks and more consistent activation of other respiratory muscles, may be typical in patients with chronic low back pain.

In healthy subjects, the diaphragm is able to perform the dual task (trunk stability and respiration) when trunk stability is challenged.¹⁹ Generally, during any body movement, with activation of the extremities during weight-bearing or weight-lifting activities and transitional movements, there is simultaneous spinal bracing and transdiaphragmatic pressure elevation.^{11,22} Intra-abdominal pressure increases, with a simultaneous decrease of intrapleural pressure, during a contraction of both the posterior (crural) and anterior (costal) portions of the diaphragm.⁷ This coordination may be compromised in patients with chronic low back pain.

Stabilizing postural activation of the diaphragm has been reported during, for example, weight lifting²⁰ and isometric activation of the extremities.²⁷ Similarly, higher inspiratory pressures and hypertrophic changes in the diaphragm have been demonstrated during exercise.^{1,6} It can be assumed that, in cases of postural instability, the insufficient stabilizers must be compensated by other muscles. A significant decrease in strength of trunk muscles, especially the extensors, in patients with low back pain has been established,^{8,23,34,37,38} suggesting that strengthening exercises of the trunk muscles may be an optimal rehabilitation strategy.

One possibility is that the lack of pos-

tural diaphragmatic activation is substituted by excessive activation of the superficial lumbar paraspinal muscles, which may lead to hypertrophy and, eventually, result in lumbar hyperlordosis and/or anterior pelvic tilt. Future research should study this mechanism as possibly contributing to or even underlying the etiology of low back pain symptoms. Furthermore, long-lasting effects and pain alleviation may be aided by achieving balanced agonist-antagonist postural activation (that is, balanced activation across the sections of the diaphragm, pelvic floor, and the abdominal wall and extensors). Research is needed to investigate the possibility that working to correct the altered function of the diaphragm specifically may contribute to alleviating low back pain symptoms by improving spinal stability.

There are several limitations to this study. First, we used a convenience sample in which the patient and control groups differed in size and some demographic characteristics. Although we used statistical methods appropriate for unbalanced designs and controlled for these differences statistically, we cannot fully exclude the possibility that these differences had some influence on the results. Second, ideally, the entire rib cage, including the whole range of diaphragm excursions, should have been imaged. Only an isolated analysis of the diaphragm was performed, which focused on the diaphragm excursions, due to the limited size of the field of view.³⁶ Although we have limited the diaphragm excursion measurements to 3 points, this is similar to studies conducted by other authors³⁶ and can be considered sufficient for this type of study. In addition, the diaphragm excursions alone may not be sufficient to understand all mechanical actions of the rib cage and related musculature. For example, individual breathing patterns may be considered along with diaphragm excursions in future research. Third, although external pressure to generate grade 4 force was applied by the same clinician (P.K.) and standardized requirements of

current MRI methodology⁴ were followed in order to reduce variation in diaphragm motion and function, force and direction were not formally assessed. Therefore, we cannot exclude the possibility that the resistance varied across the subjects.

Fourth, we did not assess the length or duration of low back pain in the patient group, only that the pain had lasted at least 6 months. Nor did we assess pain during postural tasks. However, all participants exhibited the ability to perform grade 4 force without substantial pain. Had pain been present during postural tasks, we could not have obtained readable MRI images. It could also be argued that the 4 patients who underwent a failed back surgery might have differed from the rest of the patient sample in their outcomes. Excluding these 4 patients, however, would not have substantially altered the results, although group differences in diaphragm excursion during lower and upper extremity postural tasks were somewhat (not substantially) greater, as were group differences in the correlation between total diaphragm excursion and respiratory muscle drive. Finally, poor postural function of the diaphragm may result in symptoms of low back pain and lead to chronic vertebro-genic dysfunction. However, considering all possibilities, we cannot exclude the reverse order of events. Low back pain symptoms may be indicative of an initial pathogenic insult resulting in secondary quantitative as well as qualitative adaptive changes in diaphragmatic function.

CONCLUSION

WE FOUND REDUCED DIAPHRAGM movement when isometric flexion against resistance of the upper or lower extremities was applied. The combined, more cranial position in the anterior and middle portions of the diaphragm and, particularly, the steeper slope between the middle and crural portions of the diaphragm in patients with chronic low back pain may contribute to low back pain symptoms. However,

given that the results are based on cross-sectional analysis, we cannot exclude the possibility of reverse causation. Still, the results support the theory that patients with low back pain complaints present with compromised diaphragm function, which may play an important role in postural stability. ●

KEY POINTS

FINDINGS: We found reduced diaphragm movement in patients with chronic low back pain compared to healthy controls when isometric flexion against resistance of the upper or lower extremity was applied, mainly in the anterior and middle portions. This pattern of diaphragmatic recruitment resulted in a steeper angle in the middle-posterior part of the diaphragm and likely a greater strain during activity on the ventral region of the spinal column.

IMPLICATIONS: Abnormal postural activation of the diaphragm during the postural task of isometric resistance to the extremities may serve as 1 underlying mechanism of chronic low back pain.

CAUTION: Only an isolated analysis of the diaphragm excursion was performed, due to the limited field of view. In addition, the diaphragm excursion alone may not be sufficient to understand all mechanical actions of the rib cage and related musculature. We used a convenience sample in which the patient and control groups differed in size and certain demographic characteristics. Because our study was cross-sectional in nature, we cannot exclude the possibility that low back pain symptoms may be indicative of an initial pathogenic insult resulting in secondary quantitative as well as qualitative adaptive changes in diaphragmatic function.

REFERENCES

1. Al-Bilbeisi F, McCool FD. Diaphragm recruitment during nonrespiratory activities. *Am J Respir Crit Care Med.* 2000;162:456-459.
2. American Thoracic Society. ATS/ERS statement on respiratory muscle testing. *Am J Respir Crit*

3. Aruin AS, Latash ML. Directional specificity of postural muscles in feed-forward postural reactions during fast voluntary arm movements. *Exp Brain Res.* 1995;103:323-332.
4. Chu WC, Li AM, Ng BK, et al. Dynamic magnetic resonance imaging in assessing lung volumes, chest wall, and diaphragm motions in adolescent idiopathic scoliosis versus normal controls. *Spine (Phila Pa 1976).* 2006;31:2243-2249. <http://dx.doi.org/10.1097/O1.brs.0000232822.74349.32>
5. Cresswell AG, Oddsson L, Thorstensson A. The influence of sudden perturbations on trunk muscle activity and intra-abdominal pressure while standing. *Exp Brain Res.* 1994;98:336-341.
6. DePalo VA, Parker AL, Al-Bilbeisi F, McCool FD. Respiratory muscle strength training with nonrespiratory maneuvers. *J Appl Physiol.* 2004;96:731-734. <http://dx.doi.org/10.1152/jappphysiol.00511.2003>
7. De Troyer A, Loring SH. Action of the respiratory muscles. In: Macklem PT, Mead J, eds. *Handbook of Physiology: Section 3: The Respiratory System Volume III, Parts 1 & 2: Mechanics of Breathing.* New York, NY: Oxford University Press; 1986:443-462.
8. Dvir Z, Keating JL. Trunk extension effort in patients with chronic low back dysfunction. *Spine (Phila Pa 1976).* 2003;28:685-692. <http://dx.doi.org/10.1097/O1.BRS.0000051917.04731.A4>
9. Ebenbichler GR, Oddsson LI, Kollmitzer J, Erim Z. Sensory-motor control of the lower back: implications for rehabilitation. *Med Sci Sports Exerc.* 2001;33:1889-1898.
10. Hagins M, Lamberg EM. Individuals with low back pain breathe differently than healthy individuals during a lifting task. *J Orthop Sports Phys Ther.* 2011;41:141-148. <http://dx.doi.org/10.2519/jospt.2011.3437>
11. Hemborg B, Moritz U, Lowing H. Intra-abdominal pressure and trunk muscle activity during lifting. IV. The causal factors of the intra-abdominal pressure rise. *Scand J Rehabil Med.* 1985;17:25-38.
12. Hides JA, Stokes MJ, Saide M, Jull GA, Cooper DH. Evidence of lumbar multifidus muscle wasting ipsilateral to symptoms in patients with acute/subacute low back pain. *Spine (Phila Pa 1976).* 1994;19:165-172.
13. Hodges P. Lumbopelvic stability: a functional model of biomechanics and motor control. In: Richardson C, Hodges P, Hides J, eds. *Therapeutic Exercise for Lumbopelvic Stabilization.* 2nd ed. Sydney, Australia: Churchill Livingstone; 2004:13-28.
14. Hodges PW. The role of the motor system in spinal pain: implications for rehabilitation of the athlete following lower back pain. *J Sci Med Sport.* 2000;3:243-253.
15. Hodges PW, Butler JE, McKenzie DK, Gandevia SC. Contraction of the human diaphragm during rapid postural adjustments. *J Physiol.* 1997;505 pt 2:539-548.
16. Hodges PW, Cresswell AG, Daggfeldt K, Thorstensson A. In vivo measurement of the effect of intra-abdominal pressure on the human spine. *J Biomech.* 2001;34:347-353.
17. Hodges PW, Cresswell AG, Daggfeldt K, Thorstensson A. Three dimensional preparatory trunk motion precedes asymmetrical upper limb movement. *Gait Posture.* 2000;11:92-101.
18. Hodges PW, Eriksson AE, Shirley D, Gandevia SC. Intra-abdominal pressure increases stiffness of the lumbar spine. *J Biomech.* 2005;38:1873-1880. <http://dx.doi.org/10.1016/j.jbiomech.2004.08.016>
19. Hodges PW, Gandevia SC. Activation of the human diaphragm during a repetitive postural task. *J Physiol.* 2000;522 pt 1:165-175.
20. Hodges PW, Gandevia SC. Changes in intra-abdominal pressure during postural and respiratory activation of the human diaphragm. *J Appl Physiol.* 2000;89:967-976.
21. Hodges PW, Richardson CA. Altered trunk muscle recruitment in people with low back pain with upper limb movement at different speeds. *Arch Phys Med Rehabil.* 1999;80:1005-1012.
22. Hodges PW, Richardson CA. Relationship between limb movement speed and associated contraction of the trunk muscles. *Ergonomics.* 1997;40:1220-1230. <http://dx.doi.org/10.1080/001401397187469>
23. Iwai K, Nakazato K, Irie K, Fujimoto H, Nakajima H. Trunk muscle strength and disability level of low back pain in collegiate wrestlers. *Med Sci Sports Exerc.* 2004;36:1296-1300.
24. Janssens L, Brumagne S, Polspoel K, Troosters T, McConnell A. The effect of inspiratory muscles fatigue on postural control in people with and without recurrent low back pain. *Spine (Phila Pa 1976).* 2010;35:1088-1094. <http://dx.doi.org/10.1097/BRS.0b013e3181bee5c3>
25. Kalpakcioglu B, Altinbilek T, Senel K. Determination of spondylolisthesis in low back pain by clinical evaluation. *J Back Musculoskelet Rehabil.* 2009;22:27-32. <http://dx.doi.org/10.3233/BMR-2009-0212>
26. Kendall FP, McCreary EK, Provance PG, Rodgers MM, Romani WA. *Muscles: Testing and Function with Posture and Pain.* 5th ed. Baltimore, MD: Lippincott Williams & Wilkins; 2005.
27. Kolar P, Sulc J, Kyncl M, et al. Stabilizing function of the diaphragm: dynamic MRI and synchronized spirometric assessment. *J Appl Physiol.* 2010;109:1064-1071. <http://dx.doi.org/10.1152/jappphysiol.01216.2009>
28. McGill SM. Low back stability: from formal description to issues for performance and rehabilitation. *Exerc Sport Sci Rev.* 2001;29:26-31.
29. Miller MR, Hankinson J, Brusasco V, et al. Standardisation of spirometry. *Eur Respir J.* 2005;26:319-338. <http://dx.doi.org/10.1183/09031936.05.00034805>
30. O'Sullivan PB, Beales DJ, Beetham JA, et al. Altered motor control strategies in subjects with sacroiliac joint pain during the active straight-leg-raise test. *Spine (Phila Pa 1976).* 2002;27:E1-8.

[RESEARCH REPORT]

31. Panjabi MM. The stabilizing system of the spine. Part I. Function, dysfunction, adaptation, and enhancement. *J Spinal Disord.* 1992;5:383-389; discussion 397.
32. Pel JJ, Spoor CW, Goossens RH, Pool-Goudzwaard AL. Biomechanical model study of pelvic belt influence on muscle and ligament forces. *J Biomech.* 2008;41:1878-1884. <http://dx.doi.org/10.1016/j.jbiomech.2008.04.002>
33. Quanjer PH, Tammeling GJ, Cotes JE, Pedersen OF, Peslin R, Yernault JC. Lung volumes and forced ventilatory flows. Report Working Party Standardization of Lung Function Tests, European Community for Steel and Coal. Official Statement of the European Respiratory Society. *Eur Respir J Suppl.* 1993;16:5-40.
34. Reid S, Hazard RG, Fenwick JW. Isokinetic trunk-strength deficits in people with and without low-back pain: a comparative study with consideration of effort. *J Spinal Disord.* 1991;4:68-72.
35. Tabachnick BG, Fidell LS. *Using Multivariate Statistics.* 5th ed. Boston, MA: Allyn & Bacon; 2007.
36. Takazakura R, Takahashi M, Nitta N, Murata K. Diaphragmatic motion in the sitting and supine positions: healthy subject study using a vertically open magnetic resonance system. *J Magn Reson Imaging.* 2004;19:605-609. <http://dx.doi.org/10.1002/jmri.20051>
37. Tsai YS, Sell TC, Smoliga JM, Myers JB, Learman KE, Lephart SM. A comparison of physical characteristics and swing mechanics between golfers with and without a history of low back pain. *J Orthop Sports Phys Ther.* 2010;40:430-438. <http://dx.doi.org/10.2519/jospt.2010.3152>
38. Yahia A, Jribi S, Ghroubi S, Elleuch M, Baklouti S, Habib Elleuch M. Evaluation of the posture and muscular strength of the trunk and inferior members of patients with chronic lumbar pain. *Joint Bone Spine.* 2011;78:291-297. <http://dx.doi.org/10.1016/j.jbspin.2010.09.008>



MORE INFORMATION
WWW.JOSPT.ORG

EARN CEUs With JOSPT's Read for Credit Program

JOSPT's **Read for Credit (RFC)** program invites *Journal* readers to study and analyze selected JOSPT articles and successfully complete online quizzes about them for continuing education credit. To participate in the program:

1. Go to www.jospt.org and click on **"Read for Credit"** in the left-hand navigation column that runs throughout the site or on the link in the **"Read for Credit"** box in the right-hand column of the home page.
2. Choose an article to study and when ready, click **"Take Exam"** for that article.
3. Login and pay for the quiz by credit card.
4. Take the quiz.
5. Evaluate the RFC experience and receive a personalized certificate of continuing education credits.

The RFC program offers you 2 opportunities to pass the quiz. You may review all of your answers—including the questions you missed. You receive **0.2 CEUs**, or 2 contact hours, for each quiz passed. The *Journal* website maintains a history of the quizzes you have taken and the credits and certificates you have been awarded in the **"My CEUs"** section of your **"My JOSPT"** account.

This article has been cited by:

1. Supriya G. Shah, Tenzin Choezom, G. Prabu Raja. 2019. Comparison of respiratory parameters in participants with and without chronic low back pain. *Journal of Bodywork and Movement Therapies* . [Crossref]
2. César Calvo-Lobo, Jaime Almazán-Polo, Ricardo Becerro-de-Bengoa-Vallejo, Marta Elena Losa-Iglesias, Patricia Palomo-López, David Rodríguez-Sanz, Daniel López-López. 2019. Ultrasonography comparison of diaphragm thickness and excursion between athletes with and without lumbopelvic pain. *Physical Therapy in Sport* . [Crossref]
3. Hannah Horris, Barton E. Anderson, R. Curtis Bay, Kellie C. Huxel Bliven. 2019. Clinical Breathing Mechanics Differ Based on Test and Position. *Journal of Sport Rehabilitation* 1-5. [Crossref]
4. Ginger Garner. Foundations for Yoga Practice in Rehabilitation 181-224. [Crossref]
5. Sun-Young Ha, Yun-Hee Sung. 2018. Effects of Vojta approach on diaphragm movement in children with spastic cerebral palsy. *Journal of Exercise Rehabilitation* 14:6, 1005-1009. [Crossref]
6. Nam G. Lee, Joshua (Sung) H. You, Chung H. Yi, Hye S. Jeon, Bong S. Choi, Dong R. Lee, Jae M. Park, Tae H. Lee, In T. Ryu, Hyun S. Yoon. 2018. Best Core Stabilization for Anticipatory Postural Adjustment and Falls in Hemiparetic Stroke. *Archives of Physical Medicine and Rehabilitation* 99:11, 2168-2174. [Crossref]
7. Mohammad Reza Nourbakhsh, Ashley Bluestein, Richard Power, Shaun Underwood, Leanna White. 2018. The effects of pelvic alignment versus hip joint mobility on shoulder internal rotation deficit and torque production – A blinded randomized controlled study. *Physical Therapy in Sport* 34, 164-173. [Crossref]
8. Ji-Eun Cho, Dal-Yeon Hwang, Joohee Hahn, Wan-Hee Lee. 2018. Use of real-time ultrasound imaging for biofeedback of diaphragm motion during normal breathing in healthy subjects. *Physical Therapy Rehabilitation Science* 7:3, 95-101. [Crossref]
9. Mireia Martí-Salvador, Laura Hidalgo-Moreno, Julio Doménech-Fernández, Juan Francisco Lisón, Maria Dolores Arguisuelas. 2018. Osteopathic Manipulative Treatment Including Specific Diaphragm Techniques Improves Pain and Disability in Chronic Nonspecific Low Back Pain: A Randomized Trial. *Archives of Physical Medicine and Rehabilitation* 99:9, 1720-1729. [Crossref]
10. Ya-Chen Chen, Tzu-Chien Hsiao. 2018. Towards estimation of respiratory muscle effort with respiratory inductance plethysmography signals and complementary ensemble empirical mode decomposition. *Medical & Biological Engineering & Computing* 56:7, 1293-1303. [Crossref]
11. Vikram Mohan, Aatit Paungmali, Patraporn Sitilerpisan, Umami F. Hashim, Munifa B. Mazlan, Tuan N. Nasuha. 2018. Respiratory characteristics of individuals with non-specific low back pain: A cross-sectional study. *Nursing & Health Sciences* 20:2, 224-230. [Crossref]
12. Esra Dülger, Sevil Bilgin, Elif Bulut, Deniz İnal İnce, Nezire Köse, Ceyhan Türkmen, Hatice Çetin, Jale Karakaya. 2018. The effect of stabilization exercises on diaphragm muscle thickness and movement in women with low back pain. *Journal of Back and Musculoskeletal Rehabilitation* 31:2, 323-329. [Crossref]
13. Elżbieta Szczygieł, Jędrzej Blaut, Katarzyna Zielonka-Pycka, Krzysztof Tomaszewski, Joanna Golec, Dorota Czechowska, Agata Masłoń, Edward Golec. 2018. The Impact of Deep Muscle Training on the Quality of Posture and Breathing. *Journal of Motor Behavior* 50:2, 219-227. [Crossref]
14. Vikram , Mohan, Aatit Paungmali, Patraporn Sitilerpisan, Sulaiman Md Dom, Umami Farhana Hashim, Siti Nor Binti Daud, Manikumar Muthiah. 2018. Reliability of diaphragmatic mobility assessment: A systematic review. *Polish Annals of Medicine* . [Crossref]
15. Ginger Garner. Foundations for Yoga Practice in Rehabilitation 263-307. [Crossref]
16. Marlon L. Wong, Rachael G. Anderson, Kelsey Garcia, Elissa M. Housmann, Erin McHale, Gregory S. Goldberger, Lawrence P. Cahalin. 2017. The effect of inspiratory muscle training on respiratory variables in a patient with ankylosing spondylitis: A case report. *Physiotherapy Theory and Practice* 33:10, 805-814. [Crossref]
17. Zuzana Hirjaková, Kateřina Neumannová, Jana Kimijanová, Kristína Šuttová, Miroslav Janura, František Hlavačka. 2017. Breathing changes accompanying balance improvement during biofeedback. *Neuroscience Letters* 651, 30-35. [Crossref]
18. Luca Cavaggioni. 2017. Diaphragm, Core Stability & Low Back Pain: A Rehabilitative-Preventive Perspective. *MOJ Orthopedics & Rheumatology* 7:5. . [Crossref]

19. Rylee J. Stephens, Mitchell Haas, William L. Moore, Jordan R. Emmil, Jayson A. Sipress, Alex Williams. 2017. Effects of Diaphragmatic Breathing Patterns on Balance: A Preliminary Clinical Trial. *Journal of Manipulative and Physiological Therapeutics* **40**:3, 169-175. [[Crossref](#)]
20. Mary Massery. Physical Therapy Following Phrenic Nerve Graft Surgery: Implications Far Beyond Breathing 129-137. [[Crossref](#)]
21. Nathan J. Hellyer, Nicholas M. Andreas, Andrew S. Bernstetter, Kathryn R. Cieslak, Gerad F. Donahue, Elizabeth A. Steiner, John H. Hollman, Andrea J. Boon. 2017. Comparison of Diaphragm Thickness Measurements Among Postures Via Ultrasound Imaging. *PM&R* **9**:1, 21-25. [[Crossref](#)]
22. Nele Beeckmans, Astrid Vermeersch, Roeland Lysens, Peter Van Wambeke, Nina Goossens, Tinne Thys, Simon Brumagne, Lotte Janssens. 2016. The presence of respiratory disorders in individuals with low back pain: A systematic review. *Manual Therapy* **26**, 77-86. [[Crossref](#)]
23. MASAFUMI TERADA, KYLE B. KOSIK, RYAN S. MCCANN, PHILLIP A. GRIBBLE. 2016. Diaphragm Contractility in Individuals with Chronic Ankle Instability. *Medicine & Science in Sports & Exercise* **48**:10, 2040-2045. [[Crossref](#)]
24. P. Bitnar, J. Stovicek, R. Andel, J. Arlt, M. Arltova, M. Smejkal, P. Kolar, A. Kobesova. 2016. Leg raise increases pressure in lower and upper esophageal sphincter among patients with gastroesophageal reflux disease. *Journal of Bodywork and Movement Therapies* **20**:3, 518-524. [[Crossref](#)]
25. Jeong-Hyeon Kang, Hye-Rim Suh, Chang-Yong Kim, Hyeong-Dong Kim, Hyungkun Kim. 2016. Influence of Difficulty Variation of the Core Stabilization Exercise on Thickness Changes of Abdominal Muscles in Healthy Subjects: A Pilot Study. *The Journal of Korean Physical Therapy* **28**:2, 112-118. [[Crossref](#)]
26. Hideo KANEKO, Tomomi KOBA, Risa TOKUNAGA. 2016. Differences in the Respiratory Function of Female University Students with and without Non-specific Chronic Low Back Pain. *Rigakuryoho kagaku* **31**:6, 799-804. [[Crossref](#)]
27. Salami O. Ismaila, Olutayo Aderele. Determination of safe weight limit of sandcrete block for block moulders 1-5. [[Crossref](#)]
28. LOTTE JANSSENS, ALISON K. MCCONNELL, MADELON PIJNENBURG, KURT CLAEYS, NINA GOOSSENS, ROELAND LYSENS, THIERRY TROOSTERS, SIMON BRUMAGNE. 2015. Inspiratory Muscle Training Affects Proprioceptive Use and Low Back Pain. *Medicine & Science in Sports & Exercise* **47**:1, 12-19. [[Crossref](#)]
29. John C. Hannon. 2014. Clinical rehabilitation. *Journal of Bodywork and Movement Therapies* **18**:3, 317-321. [[Crossref](#)]
30. Carlos Barroqueiro, Nuno Valente Morais. 2014. The effects of a global postural reeducation program on an adolescent handball player with isthmus spondylolisthesis. *Journal of Bodywork and Movement Therapies* **18**:2, 244-258. [[Crossref](#)]
31. Pavel Kolar, Alena Kobesova, Petra Valouchova, Petr Bitnar. Dynamic Neuromuscular Stabilization 93-98. [[Crossref](#)]
32. Laurie McLaughlin. Capnography assessment 147-153. [[Crossref](#)]
33. Pavel Kolar, Alena Kobesova, Petra Valouchova, Petr Bitnar. Dynamic Neuromuscular Stabilization 11-22. [[Crossref](#)]
34. Pavel Kolar, Alena Kobesova, Petra Valouchova, Petr Bitnar. Dynamic Neuromuscular Stabilization 163-167. [[Crossref](#)]
35. Draovitch Peter, O'Sullivan Eilish, Bryan Kelly. Hip Biomechanics During Sport . [[Citation](#)] [[Full Text](#)]
36. Caitlin J. Harper, Leili Shahgholi, Kathryn Cieslak, Nathan J. Hellyer, Jeffrey A. Strommen, Andrea J. Boon. 2013. Variability in Diaphragm Motion During Normal Breathing, Assessed With B-Mode Ultrasound. *Journal of Orthopaedic & Sports Physical Therapy* **43**:12, 927-931. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
37. Josephine Key. 2013. 'The core': Understanding it, and retraining its dysfunction. *Journal of Bodywork and Movement Therapies* **17**:4, 541-559. [[Crossref](#)]
38. Lotte Janssens, Simon Brumagne, Alison K. McConnell, Greet Hermans, Thierry Troosters, Ghislaine Gayan-Ramirez. 2013. Greater diaphragm fatigability in individuals with recurrent low back pain. *Respiratory Physiology & Neurobiology* **188**:2, 119-123. [[Crossref](#)]
39. Alena Kobesova, Pavel Kolar. 2013. Developmental kinesiology: Three levels of motor control in the assessment and treatment of the motor system. *Journal of Bodywork and Movement Therapies* . [[Crossref](#)]
40. Pavel Vostatek, Daniel Novák, Tomas Rychnovský, Šarka Rychnovská. 2013. Diaphragm Postural Function Analysis Using Magnetic Resonance Imaging. *PLoS ONE* **8**:3, e56724. [[Crossref](#)]
41. Lymperis (Perry) Koziris. 2012. Postactivation Potentiation. *Strength and Conditioning Journal* **34**:6, 75-76. [[Crossref](#)]

42. Ondrej Horacek, Jan Chlumsky, Radim Mazanec, Pavel Kolar, Ross Andel, Alena Kobesova. 2012. Pulmonary function in patients with hereditary motor and sensory neuropathy: A comparison of patients with and without spinal deformity. *Neuromuscular Disorders* 22:12, 1083-1089. [[Crossref](#)]